

SHORT COMMUNICATION

Photographic evidence of interspecies mating in geckos of the *Lepidodactylus lugubris* unisexual-bisexual complex (Squamata: Gekkonidae)

Donald W. Buden¹, Carlos Cianchini², Danko Taborosi³, Robert N. Fisher⁴, Aaron M. Bauer⁵, and Ivan Ineich⁶

¹ Division of Natural Sciences and Mathematics, College of Micronesia-FSM, P.O. Box 159, Kolonia, Pohnpei, Federated States of Micronesia 96941. E-mail: don_buden@comfsm.fm.

² P.O. Box 495, Kosrae, Federated States of Micronesia 96944. E-mail: cjcianchini@yahoo.com.

³ Island Research and Education Initiative (IREI), P.O.Box PS 303, Palikir, Pohnpei, Federated States of Micronesia 96941. E-mail: taborosi@gmail.com.

⁴ U.S. Geological Survey, Western Ecological Research Center, San Diego Field Station, 4165 Spruance Road, Suite 200, San Diego, California 92101 USA. E-mail: rfisher@usgs.gov.

⁵ Department of Biology, Villanova University, 800 Lancaster Avenue, Villanova, Pennsylvania 19085, USA. E-mail: aaron.bauer@villanova.edu.

⁶ Muséum national d'Histoire naturelle, Institut de Systématique, Évolution et Biodiversité (ISYEB), UMR 7205 (CNRS, EPHE, MNHN, UPMC), 57 rue Cuvier, F-75005 Paris, France. E-mail: ineich@mnhn.fr.

Keywords: clones, hybridization, intermating, Micronesia, Pacific Ocean, unisexual-bisexual species complex.

Palavras-chave: acasalamento heteroespecífico, clones, complexo de espécies unisexual-bissexual, hibridização, Micronésia, Oceano Pacífico.

Lepidodactylus lugubris (Duméril and Bibron, 1836) is an “all female” parthenogenetic species of gecko that is widely distributed in parts of mainland areas and on many islands of the Indian and Pacific Ocean basins (Bauer and Henle 1994, Ineich 1999, Zug 2013); it occurs extralimittally in the American tropics and subtropics (Henderson *et al.* 1976, Bauer *et al.*

2007, Lorvelec *et al.* 2011, Daza *et al.* 2012). *Lepidodactylus moestus* (Peters, 1867) is a bisexual species that occurs in Micronesia (in the western Pacific Ocean), from Palau eastward through the Federated States of Micronesia (FSM) to the Marshall Islands (Ota *et al.* 1995, Zug 2013) in sympatry with the former.

An interspecies mating between unisexual *Lepidodactylus lugubris* and a male of the bisexual *L. moestus* was photographed by Carlos Cianchini on Kosrae [Island], FSM, at 18:15 h on 22 August 2013 (Figure 1). The mating pair

Received 26 June 2014.

Accepted 29 September 2014.

Distributed December 2014.

was on a window frame inside a house at Pukusruk Wan village (05°21'01" N, 163°00'41" E, elev. 28 m a.s.l.) on the northeastern side of the island. This is the first direct evidence of mating between these two species. However, examples of "possible hybrids between *L. moestus* and *L. lugubris*" have been recorded from several disparate island groups in Micronesia (Ota *et al.* 1995). All the islands are distantly located in different directions from Kosrae and at, or near the limits of, the zone of sympatry between the two species—Koror and Ngergoi Islands, Palau, approximately 3000 km to the west, Kapingamarangi Atoll, FSM, 1026 km to the southwest, and Arno Atoll, Marshall Islands, 977 km to the east. In a broader context, differences in ploidy and dorsal pattern within populations of the *L. lugubris* complex have been attributed to presumed crosses between bisexual species and often backcrosses of asexual clonal females to bisexual males (Ineich 1987, 1988, 1992, Ineich and Ota 1992, 1993, Ota *et al.* 1995, Radtkey *et al.* 1995). Hybrid males derived from bisexual-unisexual crosses are rare; they exhibit gonadal and spermatogenic abnormalities and are presumed to be sterile (Ineich and Ota 1992, Yamashiro and Ota 1998). The anatomy of offspring resulting from unisexual-bisexual crosses has been studied in French Polynesia. The genital tract of the hybrid is aberrant and does not allow reproduction (Saint Girons and Ineich 1992).

Molecular and morphological studies indicate that examples of *Lepidodactylus lugubris* previously considered bisexual and having a functional genital tract were misidentified. These are cryptic, bisexual species that include *L. moestus* and at least one undescribed species, which are the respective maternal and paternal ancestors of some clonal lineages of *L. lugubris* *sensu stricto* (Volobouev *et al.* 1993, Radtkey *et al.* 1995). At least five major clones of *L. lugubris* differing in ploidy and coloration (dorsal pattern) were first recognized by Ineich (1988; see also Ineich 1999). The diploid lineages are thought to be derived from




Figure 1. Interspecies mating between *Lepidodactylus moestus* (left) and *L. lugubris* (right) photographed on a window frame inside a house on Kosrae, Federated States of Micronesia; dark area at the top of the image is the bottom edge of the window, looking out.

hybridization between bisexual species. The triploids might have resulted from backcrosses of asexual diploid clonal females with males of one of the two bisexual species, and perhaps also with other and now extinct bisexual species (Ineich 1988, 1999, Moritz *et al.* 1993, Boissinot *et al.* 1997). The *L. lugubris* in the interspecies pair photographed on Kosrae exemplifies diploid Clone-A, which is distinguished by having seven (but occasionally 6 or 8) pairs of small dark spots straddling the dorsal midline from the neck to the base of the tail (Ineich 1988, 1999). It is the most widespread of the *L. lugubris* clones (Ineich 1999, Short and Petren 2008) and has been implicated in the probable displacement and depletion of other clones and bisexual species in its spread throughout the Pacific after World War II (Ineich 1999). Now, it appears to be expanding its range into the Caribbean

(Atlantic Ocean) Basin (Lorvelec *et al.* 2011) and mainland South America (Bauer *et al.* 2007) via human introduction.

The distribution and relative abundance of the different clones and bisexual species of the *Lepidodactylus lugubris* complex within the FSM is poorly known and in need of further study. Ineich's (1999) review of variation from the entire distributional range of the unisexual-bisexual complex included relatively few specimens from the FSM. Three were from Kosrae (1 diploid Clone-A, 1 triploid Clone-B, and a male identified as likely being *L. moestus*) and several sterile individuals were from Pohnpei State, which Ineich (1999) thought to have originated from crosses between Clone-A females and *L. moestus* males. Nine other specimens were from Yap State, which included eight *L. moestus* from Yap proper and a Clone-B from one of the outer atolls. More recently, *L. lugubris*, *L. moestus*, and *L. sp.* were recorded in sympatry in Yap State on Ngulu (Buden 2010), and Sorol (Buden 2013) Atolls, in Chuuk State on Satawan (Buden 2007) and Namonuito (Buden and Taborosi in prep.) Atolls, and on Houk Island (Buden and Taborosi in prep.), but individual clones were not identified according to Ineich's (1988) clonal categories.

Additionally, Ineich (1987) observed occasional pseudocopulations between distinct clones (diploid and triploid) of *Lepidodactylus lugubris* in French Polynesia and such behavior has since been recorded in other places (McCoid and Hensley 1991, Brown and O'Brien 1993). This behavior seems to be common among unisexual lizards (Cole and Townsend 1983). Thus it seems that clonal females can copulate or pseudocopulate with other distinct clones, with one of their parental bisexual species, or with some other bisexual species within the *L. lugubris* unisexual-bisexual complex.

Acknowledgments.—We thank Steven Poe and an anonymous reviewer for their constructive comments on the manuscript. 

References

- Bauer, A. M. and K. Henle. 1994. *Das Tierreich 109, Family Gekkonidae (Reptilia, Sauria). Part 1. Australia and Oceania*. Berlin and New York. Walter De Gruyter. 306 pp.
- Bauer, A. M., T. R. Jackman, E. Greenbaum, and T. J. Papenfuss. 2007. First record of *Lepidodactylus lugubris* in Suriname. *Applied Herpetology* 4: 84–85.
- Boissinot, S., I. Ineich, L. Thaler, and C.-P. Guillaume. 1997. Hybrid origin and clonal diversity in the parthenogenetic gecko *Lepidodactylus lugubris* in French Polynesia. *Journal of Herpetology* 31: 295–298.
- Brown, S. G. and J. O'Brien. 1993. Pseudosexual and dominance behavior: Their relationship to fecundity in the unisexual gecko, *Lepidodactylus lugubris*. *Journal of Zoology* 2331: 61–69.
- Buden, D. W. 2007. Reptiles of Satawan Atoll and the Mortlock Islands, Chuuk State, Federated States of Micronesia. *Pacific Science* 61: 415–428.
- Buden, D. W. 2010. Reptiles of Ngulu Atoll, Yap State, Federated States of Micronesia. *Pacific Science* 64: 473–480.
- Buden, D. W. 2013. The reptiles of Sorol Atoll, Yap, Federated States of Micronesia. *Pacific Science* 67: 119–128.
- Cole, C. J. and C. R. Townsend. 1983. Sexual behaviour in unisexual lizards. *Animal Behaviour* 31: 724–728.
- Daza, J. D., S. L. Travers, and A. M. Bauer. 2012. New records of the mourning gecko *Lepidodactylus lugubris* (Duméril and Bibron, 1836) (Squamata: Gekkonidae) from Colombia. *Check List* 8: 164–167.
- Henderson, R. W., J. Villa, and A. M. Bauer. 1976. *Lepidodactylus lugubris* (Reptilia: Gekkonidae): a recent addition to the herpetofauna of Nicaragua. *Herpetological Review* 7: 173.
- Ineich, I. 1987. Recherches sur le peuplement et l'évolution des Reptiles terrestres de Polynésie française. Unpublished Ph.D. Thesis, Montpellier University, Université des Sciences et Techniques du Languedoc. 515 pp.
- Ineich, I. 1988. Mise en évidence d'un complexe unisexe-bisexe chez le gecko *Lepidodactylus lugubris* (Sauria, Lacertilia) en Polynésie française. *Comptes Rendus de l'Académie des Sciences. Série III* 307: 271–277.
- Ineich, I. 1992. La parthénogénèse chez les Gekkonidae (Reptilia, Lacertilia): Origine et évolution. *Bulletin de la Société zoologique de France* 117: 253–266.

- Ineich, I. 1999. Spatio-temporal analysis of the unisexual-bisexual *Lepidodactylus lugubris* complex (Reptilia Gekkonidae). Pp. 199–228 in H. Ota (ed.), *Tropical Island Herpetofauna: Origin, Current Diversity, and Conservation*. Amsterdam: Elsevier.
- Ineich, I. and H. Ota. 1992. Additional remarks on the unisexual-bisexual complex of the gecko *Lepidodactylus lugubris*, in Takapoto Atoll, French Polynesia. *Bulletin of the College of Science, University of the Ryukyus* 53: 31–39.
- Ineich, I. and H. Ota 1993. Morphological variation and distribution of the unisexual-bisexual complex of the gecko, *Lepidodactylus lugubris*, in French Polynesia and Easter Island. *Bulletin of the College of Science, University of the Ryukyus* 56: 113–120.
- Lorvelec, O., A. Levesque, and A. M. Bauer. 2011. First record of the Mourning Gecko (*Lepidodactylus lugubris*) on Guadeloupe, French West Indies. *Herpetology Notes* 4: 291–294.
- McCoid, M. J. and R. A. Hensley 1991. Pseudocopulation in *Lepidodactylus lugubris*. *Herpetological Review* 22: 8–9.
- Moritz, C., D. T. Bolger, S. C. Donnellan, and T. J. Case. 1993. Genetic diversity and the history of Pacific island geckos (*Hemidactylus* and *Lepidodactylus*). *Biological Journal of the Linnean Society* 48: 113–133.
- Ota, H., R. N. Fisher, I. Ineich, and T. J. Case. 1995. Geckos of the genus *Lepidodactylus* (Squamata: Reptilia) in Micronesia: description of a new species and reevaluation of the status of *Gecko moestus* Peters, 1867. *Copeia* 1995: 183–195.
- Radtkey, R. R., S. C. Donnellan, R. N. Fisher, C. Moritz, K. A. Hanley, and T. J. Case. 1995. When species collide: the origin and spread of an asexual species of gecko. *Proceedings of the Royal Society of London B: Biological Sciences* 259: 145–152.
- Saint Girons, H. and I. Ineich 1992. Histology of the reproductive tract of hybrids between gonochoristic males and parthenogenetic females of the *Lepidodactylus lugubris* (Duméril and Bibron, 1836) in French Polynesia (Reptilia, Gekkonidae). *Journal of Morphology* 212: 55–64.
- Short, K. H. and K. Petren. 2008. Boldness underlies foraging success of invasive *Lepidodactylus lugubris* geckos in the human landscape. *Animal Behaviour* 76: 429–437.
- Volobouev, V., G. Pasteur, I. Ineich, and B. Dutrillaux. 1993. Chromosomal evidence for a hybrid origin of diploid parthenogenetic females from the unisexual-bisexual *Lepidodactylus lugubris* complex (Reptilia: Gekkonidae). *Cytogenetics and Cell Genetics* 63: 194–199.
- Yamashiro, S. and H. Ota. 1998. Discovery of a male phenotype of the parthenogenetic gecko, *Lepidodactylus lugubris*, on Ishigakijima Island of the Yaeyama group, Ryukyu Archipelago. *Japanese Journal of Herpetology* 17: 1523–155.
- Zug, G. 2013. *Reptiles and Amphibians of the Pacific Islands: A Comprehensive Guide*. Berkeley. University of California Press. 306 pp.

Editor: Steven Poe